arXiv:1603.06551v1 [math.GR] 21 Mar 2016

Finite soluble groups with nilpotent wide subgroups

V. S. Monakhov and I. L. Sokhor

March 22, 2016

Abstract

A subgroup H of a finite group G is wide if each prime divisor of the order of G divides the order of H. We obtain the description of finite soluble groups with no wide subgroups. We also prove that a finite soluble group with nilpotent wide subgroups has the quotient group by its hypercenter with no wide subgroups.

Keywords: finite groups, soluble groups, nilpotent groups.

MSC2010: 20D20, 20E28, 20E34.

1 Introduction

All groups in this paper are finite. All notations and terminology are standard. The reader is referred to [1, 2] if necessary.

Let G be a group. We use $\pi(G)$ to denote the set of all prime devisors of $|G|$. By $|\pi(G)|$ we denote a number of different prime devisors of $|G|$. A subgroup H of a group G is said to be wide if $\pi(H) = \pi(G)$.

Let k be a positive integer. A group G is called k-primary if $|\pi(G)| = k$. If $|\pi(G)| = 1$ or $|\pi(G)| = 2$, then G is said to be primary or biprimary, respectively. A group G is quasi-k-primary if $|\pi(G)| > k$ and $|\pi(M)| \leq k$ for every maximal subgroup M of G. Obviously, quasi-k-primary groups have no wide subgroups.

A quasi-1-primary group is also called quasiprimary, and a quasi-2-primary group is also called quasibiprimary [3]. It is clear that the order of a nilpotent quasiprimary group is equal to pq, where p and q are primes. A nonnilpotent quasiprimary group G can be represented as the semidirect product $G = [E_{p^a}]Z_q$ of a normal elementary abelian group E_{p^a}, $|E_{p^a}| = p^a$, by a cyclic group Z_q, $|Z_q| = q$, where a is the exponent of p. This follows from Schmidt theorem [4] on groups all of whose subgroups are nilpotent.

S. S. Levischenko investigated quasibiprimary groups [3]. A soluble quasibiprimary group G can be represented as the semidirect product $[P]M$ of an elementary abelian Sylow p-subgroup P and a quasiprimary subgroup M, which is a maximal subgroup of G [3, Theorem 3.1]. In an insoluble quasibiprimary group G the Frattini subgroup $\Phi(G)$ is primary [3, Theorem 2.2], the quotient group $G/\Phi(G)$ is a simple group, and all such simple groups are enumerated [3, Theorem 2.1].

It is natural to study the structure of quasi-k-primary groups for any positive integer k. Most of simple groups is quasi-k-primary. Simple groups that are not quasi-k-primary are enumerated in [5, 3.8]. It is satisfied the question of V. S. Monakhov in Kourovka notebook [6, 11.64].

We obtain the description of soluble quasi-k-primary groups and soluble groups with nilpotent wide subgroups.

2 Preliminaries

Lemma 1. *A soluble quasi-k-primary group is $(k+1)$-primary.*

Proof. Let G be a soluble quasi-k-primary group and M be a maximal subgroup of G. Then $|\pi(G)| \geq k+1$ and $|\pi(M)| \leq k$. In a soluble group maximal subgroups have prime indices. Therefore

$$|G : M| = p^{\alpha},\ p \in \pi(G),\ \alpha \in \mathbb{N}.$$

Since $|G| = |M| \cdot |G : M|$, we have $|\pi(G)| \leq k+1$, and $|\pi(G)| = k+1$. Lemma 1 is proved.

Lemma 2. *If G is a soluble quasi-k-primary group and N is its normal Hall subgroup, then G/N is a quasi-l-primary group, where $l = k - |\pi(N)|$.*

Proof. By Lemma 1, G is $(k+1)$-primary, hence

$$|\pi(G/N)| = |\pi(G)| - |\pi(N)| = k + 1 - |\pi(N)| = l + 1.$$

Let M/N be a maximal subgroup of G/N. Then M is a maximal subgroup of G. Therefore $|\pi(M)| < |\pi(G)|$. As N is a Hall subgroup of G, we obtain

$$|\pi(M/N)| = |\pi(M)| - |\pi(N)| < |\pi(G)| - |\pi(N)| = |\pi(G/N)| = l + 1.$$

Thus G/N is $(l+1)$-primary, and any maximal subgroup of G is not more than l-primary. Consequently, G/N is quasi-l-primary. Lemma 2 is proved.

Lemma 3. ([2, IV.2.11]) *If all Sylow subgroups of a group G are cyclic, then the derived subgroup G' is a cyclic Hall subgroup and the quotient group G/G' is cyclic.*

3 The structure of soluble quasi-k-primary groups

Theorem 1. *Let G be a soluble group. Then the following statements are equivalent.*

(1) *G is a quasi-k-primary group.*

(2) *Every normal subgroup of G is a Hall subgroup.*

(3) *Every maximal subgroup of G is a Hall subgroup.*

(4) *$G = [N]M$, where N is a minimal normal and Sylow subgroup of G, M is a quasi-$(k-1)$-primary and maximal subgroup.*

Proof. Check that (1) implies (2). Let G be a soluble quasi-k-primary group and N be a normal subgroup of G, $\tau = \pi(G) \setminus \pi(N)$. Suppose that N is not a Hall subgroup of G. Since G is a quasi-k-primary group and N is its proper subgroup, we have $|\pi(N)| \leq k$ and $\tau \neq \emptyset$. As G is a soluble group, in G there is τ-Hall subgroup M. Now, $(|M|, |N|) = 1$, so $M \cap N = 1$ and $NM = [N]M < G$. But

$$\pi([N]M) = \pi(N) \cup \pi(M) = \pi(N) \cup \tau = \pi(G),\ |\pi(G)| = k + 1,$$

that is, in a quasi-k-primary group G there is a proper $(k+1)$-primary subgroup $[N]M$, a contradiction. Thus N is a Hall subgroup of G.

Now we prove that (3) follows from (2). Assume that in a soluble group G every normal subgroup is a Hall subgroup. Let M be a maximal subgroup and N a minimal normal subgroup of G. Then N is a Sylow subgroup of G. If N does not belong to M, then $G = NM$. And since N is abelian, $N \cap M = 1$ and subgroup M is a Hall subgroup. Let $N \subseteq M$. Then M/N is a maximal subgroup of G/N. It is clear that every normal subgroup of G/N is a Hall subgroup. Then by induction M/N is a Hall subgroup of G/N. Therefore M is also a Hall subgroup of G.

Check that (3) implies (1). Assume that every maximal subgroup of a soluble group G is a Hall subgroup. Let M be a maximal subgroup of G and $|\pi(G)| = k + 1$. Since M is a Hall subgroup, we have $|\pi(M)| = |\pi(G)| - 1 = k$, and G is a quasi-k-primary group.

Thus, (1), (2) and (3) are equivalent.

Check that (1) implies (4). Let G be a soluble quasi-k-primary group and N its minimal normal subgroup. In view of (2), N is a Sylow p-subgroup of G. By SchurZassenhaus theorem [1, Theorem 4.32], in G there is a subgroup M such that $G = [N]M$. Applying Lemma 1, G is $(k+1)$-primary. Hence M is k-primary. Suppose that in M there is a proper k-primary subgroup M_1. Then in G there is a proper $(k+1)$-primary subgroup $[N]M_1$, a contradiction. Consequently, M is quasi-$(k-1)$-primary.

Finely we prove that (1) follows from (4). Let $G = [N]M$, where N is a minimal normal and Sylow subgroup of G, M is its quasi-$(k-1)$-primary and maximal subgroup. Then by

Lemma 1,

$$|\pi(M)| = k,\ |\pi(G)| = |\pi([N]M)| = |\pi(N)| + |\pi(M)| = k + 1.$$

Show that every maximal subgroup K of G is not more than k-primary. If $NK = G$, then $N \cap K = 1$ and $K \simeq M$. Since M is a quasi-$(k-1)$-primary soluble subgroup of G, using Lemma 1, we obtain that M is k-primary. And so K is also a k-primary group. Assume that NK is a proper subgroup of G. Then $N \subseteq K$ and $K = N(K \cap M)$ by Dedekind identity. As M is quasi-$(k-1)$-primary, we obtain that $L \cap M$ is not more than $(k-1)$-primary. Hence $K = [N](K \cap M)$ is not more than k-primary. Thus (4) and (1) are equivalent. Theorem 1 is proved.

Note that π-soluble groups with certain Hall maximal subgroups are investigated in [7].

Corollary 1.1. *In a soluble quasi-k-primary group the Frattini subgroup is trivial.*

Proof. The Frattini subgroup of any group can never be a Hall subgroup [1, Theorem 4.33]. It remains only to use Statement (2) of Theorem 1.

Corollary 1.2. *In a soluble quasi-k-primary group the Fitting subgroup is a Hall subgroup and every its Sylow subgroup is a minimal normal subgroup.*

Proof. By Theorem 1 (2) the Fitting subgroup of a soluble quasi-k-primary group G is a Hall subgroup. And so every its Sylow subgroup P is a Sylow subgroup of G. Moreover, since the Fitting subgroup is nilpotent, P is a characteristic subgroup. Consequently, P is normal in G. At the same time by Theorem 1 (2), a minimal normal subgroup of a soluble quasi-k-primary group is a Sylow subgroup. Corollary 1.2 is proved.

Corollary 1.3. *Let N be a normal subgroup of a group G. If G is a soluble quasi-k-primary group, then the quotient group G/N is a soluble quasi-l-primary group, where $l = k - |\pi(N)|$.*

Proof. It follows from Theorem 1 (2) and Lemma 2.

If a group G has a normal series whose factors are isomorphic to Sylow subgroups, then we say G has a Sylow tower.

Corollary 1.4. *A soluble quasi-k-primary group has a Sylow tower.*

Proof. We use induction on k. Let G be a soluble quasi-k-primary group. Then by Statement (2) and (4) of Theorem 1, G can be represented as $G = [P_1]M_1$, where P_1 is a minimal normal and Sylow subgroup of G, M_1 is a quasi-$(k-1)$-primary and maximal subgroup. By the induction hypothesis, M_1 has a Sylow tower. Hence G also has a Sylow tower. Corollary 1.4 is proved.

A positive integer n is said to be squarefree if p^2 does not divide n for all primes p. A group is supersoluble if all its chief factors are of prime orders.

Corollary 1.5. *The order of a group G is squarefree if and only if G is a supersoluble quasi-k-primary group, where $k = |\pi(G)| - 1$. In particular, a supersoluble quasi-k-primary group is metacyclic.*

Proof. Let the order of a group G be squarefree and $k = |\pi(G)| - 1$. Then all Sylow subgroups of G are cyclic and G is supersoluble and metacyclic by Lemma 3. It is clear that $|\pi(X)| < |\pi(G)|$ for every proper subgroup X of G, that is, G is quasi-k-primary.

Converse, let G be a supersoluble quasi-k-primary group. Apply induction on k. By Theorem 1 (4), $G = [N]M$, where N is a minimal normal and Sylow subgroup of G, M is a quasi-$(k-1)$-primary and maximal subgroup. In view of [1, Theorem 4.48], a minimal normal subgroup of a supersoluble group is of prime order, hence $|N| = p$, $p \in \pi(G)$. A subgroup M is supersoluble and quasi-$(k-1)$-primary. By the induction hypotheses, the order of M is squarefree. Hence the order of G is also squarefree. And so G is metacyclic by Lemma 3. Corollary 1.5 is proved.

Corollary 1.6. *The derived length of a soluble quasi-k-primary group G does not exceed*

$$\min\{|\pi(G)|,\ \max\{1 + a_i \mid i = 1, 2, \ldots, t\}\},$$

where $|F(G)| = p_1^{a_1} p_2^{a_2} \ldots p_t^{a_t}$.

Proof. Let $d(G)$ and $r(G)$ be the derived length and the rank of a soluble quasi-k-primary group G, respectively. By Theorem 1 (4), all Sylow subgroups of G are elementary abelian and the length of chief series of G equals $|\pi(G)|$. Hence $d(G) \leq |\pi(G)|$. Soluble groups with abelian Sylow subgroups are considered in paragraph VI.14 [2]. Therefore all statements applying to A-groups (soluble groups with abelian Sylow subgroups) of this paragraph is correct for quasi-k-primary groups. In particular, $d(G) \leq 1 + r(G)$ by [2, VI.14.18], and $r(G) \leq u$ by [2, VI.14.31], where u is the maximal number of generating Sylow subgroups in the Fitting subgroup $F(G)$. Since Sylow subgroups of G are elementary abelian, we have $u = \max\{a_i \mid i = 1, \ldots, t\}$. It follows that

$$d(G) \leq \min\{|\pi(G)|, \max\{1 + a_i \mid i = 1, \ldots, t\}\}.$$

Corollary 1.6 is proved.

Substituting $k = 2$ in Theorem 1, we obtain the result of S. S. Levischenko.

Corollary 1.7. ([3, Theorem 3.1]) *A soluble group G is quasibiprimary if and only if G is equal to the semidirect product $[P]M$ of its elementary abelian Sylow p-subgroup P and quasiprimary subgroup M, which is also a maximal subgroup of G.*

4 Soluble groups with nilpotent wide subgroups

Let G be a nontrivial group, $Z_0(G) = 1$, $Z_1(G) = Z(G)$, $Z_2(G)/Z_1(G) = Z(G/Z_1(G))$, ..., $Z_i(G)/Z_{i-1}(G) = Z(G/Z_{i-1}(G))$, ... Then the subgroup $Z_\infty(G) = \bigcup_{i=0}^{\infty} Z_i(G)$ is called the hypercenter of G.

Obviously, $Z(G/Z_\infty(G)) = 1$.

Theorem 2. *If in a soluble group G every maximal subgroup M such that $\pi(M) = \pi(G)$ is nilpotent, then the quotient group $G/Z_\infty(G)$ is quasi-k-primary, where $k = |\pi(G/Z_\infty(G))| - 1$.*

Proof. Suppose that the quotient group $\overline{G} = G/Z_\infty(G)$ is not quasi-k-primary, where $k = |\pi(G/Z_\infty(G))| - 1$. Then in $\overline{G}$ there is a maximal subgroup $\overline{M} = M/Z_\infty(G)$ such that $\pi(\overline{M}) = \pi(\overline{G})$. Hence M is a maximal subgroup of G and $\pi(M) = \pi(G)$. By the hypotheses, M is nilpotent. Therefore $\overline{M}$ is also nilpotent. Since $\overline{M}$ is a maximal subgroup of a soluble group $\overline{G}$, we obtain that $|\overline{G} : \overline{M}| = p^\alpha$ for some $p \in \pi(G)$ and positive integer α. Let $\overline{M}_p$ be a Sylow p-subgroup of $\overline{M}$, $\overline{G}_p$ be a Sylow p-subgroup of $\overline{G}$ containing $\overline{M}_p$. Then $\overline{M}_p$ is a proper subgroup of $\overline{G}_p$. Therefore $\overline{M}_p$ is normal in $\overline{G}$. Now there is a nontrivial element $\overline{x}$ of $\overline{M}_p \cap Z(\overline{G}_p)$ such that it belongs to the center of $\overline{G}$. This contradicts the fact that $Z(G/Z_\infty(G)) = 1$. Theorem 2 is proved.

Corollary 2.1. *If all wide subgroups of a group G are nilpotent, then $G/Z_\infty(G)$ is quasi-k-primary, where $k = |\pi(G/Z_\infty(G))| - 1$.*

The converse of Theorem 2 is not true. For example, let $G = S_3 \times Z_6$. Here S_3 is the symmetric group of order 3, Z_6 is the cyclic group of order 6. Then $Z_\infty(G) = Z_6$, and so $G/Z_\infty(G) \simeq S_3$ is quasiprimary, but the wide maximal subgroup $M = S_3 \times Z_2$ is nonnilpotent.

The following question becomes natural.

What is the structure of a finite soluble group all of whose wide subgroups are supersoluble?

References

[1] V. S. Monakhov, *Introduction to the Theory of Finite groups and their Classes*, Vyshejshaja shkola, Minsk, 2006 (In Russia).

[2] B. Huppert, *Endliche Gruppen I*, Berlin-Heidelberg-New York, 1967.

[3] S. S. Lewischenko, Finite quasibiprimary groups, *Groups defined by properties of group systems* : collection of scientific papers, Inst. matem. AN USSR, Kiev (1979), 83–97 (In Russia).

[4] O. Yu. Schmidt, *Groups all of whose subgroups are special*, Mat. Sb. **31** (1924), 366–372.

[5] Q. Zhang, L. Wang, Finite non-abelian simple groups which contain a non-trivial semipermutable subgroup, *Algebra Colloquium*, **12** (2005), 301–307.

[6] *The Kourovka Notebook: Unsolved Problems in Group Theory*, Institute of Mathematics, Russian Academy of Sciences, Novosibirsk, 2014.

[7] V. S. Monakhov, Finite π-solvable groups whose maximal subgroups have the Hall property, *Math. Notes*, **84** (2008), P. 363–366.